

Scotland's Rural College

Determining the variation in premaxillary and dentary morphology that may underlie beak shape between two pure layer lines

Struthers, S; Andersson, Bjorn; Schmutz, Matthias; McCormack, Heather; Wilson, Peter; Dunn, Ian; Sandilands, V; Schoenebeck, Jeffrey

Published in:
Poultry Science

DOI:
[10.1016/j.psj.2021.101500](https://doi.org/10.1016/j.psj.2021.101500)

Print publication: 01/12/2021

Document Version

Version created as part of publication process; publisher's layout; not normally made publicly available

[Link to publication](#)

Citation for pulished version (APA):

Struthers, S., Andersson, B., Schmutz, M., McCormack, H., Wilson, P., Dunn, I., Sandilands, V., & Schoenebeck, J. (2021). Determining the variation in premaxillary and dentary morphology that may underlie beak shape between two pure layer lines. *Poultry Science*, 100(12), [101500].
<https://doi.org/10.1016/j.psj.2021.101500>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Journal Pre-proof

Determining the variation in premaxillary and dentary bone morphology that may underlie beak shape between two pure layer lines

S. Struthers , B. Andersson , M. Schmutz , H.A. McCormack ,
P.W. Wilson , I.C. Dunn , V. Sandilands , J.J. Schoenebeck

PII: S0032-5791(21)00522-8
DOI: <https://doi.org/10.1016/j.psj.2021.101500>
Reference: PSJ 101500



To appear in: *Poultry Science*

Received date: 10 June 2021
Accepted date: 14 September 2021

Please cite this article as: S. Struthers , B. Andersson , M. Schmutz , H.A. McCormack , P.W. Wilson , I.C. Dunn , V. Sandilands , J.J. Schoenebeck , Determining the variation in premaxillary and dentary bone morphology that may underlie beak shape between two pure layer lines, *Poultry Science* (2021), doi: <https://doi.org/10.1016/j.psj.2021.101500>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Inc. on behalf of Poultry Science Association Inc.
This is an open access article under the CC BY-NC-ND license
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

BEAK BONE MORPHOLOGY IN PURE LINE LAYING HENS

**Determining the variation in premaxillary and dentary bone morphology that may
underlie beak shape between two pure layer lines**

S. Struthers^{*,#,1}, B. Andersson[§], M. Schmutz[§], H. A. McCormack^{*}, P. W. Wilson^{*}, I. C.
Dunn^{*}, V. Sandilands[#], and J. J. Schoenebeck^{*}

^{*}The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of
Edinburgh, Edinburgh, UK EH25 9RG; [#]Department of Agriculture, Horticulture, and
Engineering Science, Scotland's Rural College, Edinburgh, UK EH25 9RG; [§]Lohmann
Breeders GmbH, Cuxhaven, DE 27472

¹Corresponding author: sarah.struthers@ed.ac.uk

Sarah Struthers

The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh,
Edinburgh, UK EH25 9RG

Scientific Section: Management and Production

ABSTRACT

Beak treatment is an effective method of reducing the damage inflicted by severe feather pecking (SFP) but there is significant pressure to eliminate these treatments and rely solely on alternative strategies. Substantial variation in beak shape exists within non-beak treated layer flocks and beak shape appears to be heritable. There is the potential to use this pre-existing variation and genetically select for hens whose beak shapes are less apt to cause damage during SFP. To do this, we must first understand the range of phenotypes that exist for both the external beak shape and the bones that provide its structure. The objective of this study was to determine the variation in premaxillary (within the top beak) and dentary (within the bottom beak) bone morphology that exists in two non-beak treated pure White Leghorn layer lines using geometric morphometrics to analyze radiographs. Lateral head radiographs were taken of 825 hens and the premaxillary and dentary bones were landmarked. Landmark coordinates were standardized by Procrustes superimposition and the covariation was analyzed by principal components analysis and multivariate regression using Geomorph (an R package). Three principal components (PCs) explained 85% of total premaxillary bone shape variation and showed that the shape ranged from long and narrow with pointed bone tips to short and wide with more curved tips. Two PCs explained 81% of total dentary bone shape variation. PC1 described the dentary bone length and width and PC2 explained the angle between the bone tip and its articular process. For both bones, shape was significantly associated with bone size and differed significantly between the two lines. Bone size accounted for 42% of the total shape variation for both bones. Together, the results showed a range of phenotypic variation in premaxillary and dentary bone shape, which in turn may influence beak shape. These bone phenotypes will guide further quantitative genetic and

behavioral analyses that will help identify which beaks shapes cause the least damage when birds engage in SFP.

Key words: Lohmann, principal components analysis, radiography, bone shape, laying hen

INTRODUCTION

Animal welfare and how food production animals are raised and managed are becoming more important to consumers and society in general (Napolitano et al., 2010; Spooner et al., 2014; Dunne and Sietou, 2020). This concern for animal welfare has led to widespread legislation regarding how animals are housed (Centner, 2010). In the egg production industry, the use of alternative housing systems such as free-range or whole-barn housing has increased, largely in response to the shift away from and/or ban on cages (either conventional or furnished) in many countries. For example, 56% of the 11 billion eggs produced in the United Kingdom (UK) in 2019 came from free-range systems (British Egg Industry Council, 2019). In the United States, over 29% of laying hens are housed in cage-free systems, marking a 15% increase since 2016 (United Egg Producers, 2021). These alternative systems allow birds more space and the ability to express their full behavioral repertoire (Muir et al., 2014). However, the large group sizes and unstable social hierarchies in these systems can contribute to outbreaks of severe feather pecking (SFP) (Bilčík and Keeling, 2000; Lay et al., 2011; Muir et al., 2014). SFP is the pecking at, pulling, and removal of feathers which can cause damage to the integument of the birds, often resulting in wounds and cannibalism (Savory, 1995). The behavior is a serious welfare and economic concern as the loss of feathers can result not only in pain and mortality but in increased feed costs and poor feed efficiency (Leeson and Walsh, 2004).

The most effective method to control SFP is beak treatment, which both blunts and shortens beak length. Whether by hot blade trimming or infrared methods, beak treatment raises its own welfare concerns. Hot blade trimming results in acute pain and may cause neuroma formation and chronic pain, depending on the age of the bird at trimming and the severity of the trim (Lunam et al., 1996; Gentle et al., 1997). Infrared beak treatment has much less of a negative impact on bird welfare than hot blade trimming (Gentle and McKeegan, 2007; Dennis et al., 2009; Struthers et al., 2019) but concern regarding any form of beak manipulation has led to many national governments banning these treatments or, such as in the UK, allowing them only until better SFP prevention methods are found (Department for Environment Food and Rural Affairs, 2010; Scottish Executive, 2010).

Alternative management practices to beak treatment have been studied and led to some successes in reducing the incidence of SFP; however, the behavior remains unpredictable and difficult to control (Jendral and Robinson, 2004). More recently, there has been an increased focus on selective breeding against SFP (Rodenburg et al., 2003; Bennewitz et al., 2014; Grams et al., 2015). The incorporation of meaningful behavioral data into a breeding program can be challenging; therefore, quantifiable outcomes such as plumage cover, liveability, and beak shape are measured (Ellen et al., 2019). Plumage cover has been reported to have a moderate heritability and selection for the trait has reduced the incidence of SFP (Brinker et al., 2014; Icken et al., 2017). However, a disadvantage is that only the victims of feather pecking can be detected and it is time consuming to measure (Brinker et al., 2014). A hen's liveability in relation to feather pecking depends both on her ability to avoid being feather pecked and the tendency of her flock mates to feather peck (Ellen and Bijma, 2019). Group selection for liveability reduced beak-related injuries and mortality in non-beak treated laying hens (Kuo et al., 1991; Craig and Muir, 1993; Rodenburg et al., 2010; Ellen and Bijma, 2019) but continual selection caused heritability to

decrease over time meaning that further improvement becomes more and more difficult (Ellen and Bijma, 2019).

Beak shape differs between bird species because each species has evolved a specific shape to fit both their diet and environmental niche (Grant and Grant, 1993). The beak is a highly specialized and complex organ that serves many important functions including feeding, drinking, grooming, parasite removal, and defense (Lunam, 2005). In chickens, the natural shape of the beak resembles that of a hook, with the sharp tip of the top beak extending over the bottom beak. The internal structure of the top and bottom beak are provided by the premaxillary and dentary bones, respectively (Lunam, 2005). To what extent beak shape can contribute towards reducing the incidence of SFP is still not fully understood. It is also not well understood if phenotypic variation in beak shape is due primarily to genetics, environmental factors, or a combination of both. Substantial variation in beak shape exists within non-beak treated layer flocks and beak shape appears to be heritable (Icken et al., 2017). Using a device that measured the difference in length between the top and bottom beak (i.e., top beak overhang), Icken et al. (2017) found that when hens had naturally shorter (blunter) top beaks, mortality was reduced and plumage cover improved. This suggests that there is the potential to use this pre-existing variation and selectively breed hens whose beak shapes are less apt to cause damage during SFP.

Beak morphology in poultry has previously been described using linear measurements such as length, width, and depth (Marchant-Forde et al., 2008; Carruthers et al., 2012; McKeegan and Philbey, 2012). However, these measurements help highlight differences in size rather than true shape differences. Interpretation of these measurements is also difficult when the study specimen occupies three-dimensional space. Geometric morphometrics is the analysis of morphological shape using landmark coordinates rather than linear measurements

(Zelditch et al., 2012). Geometric morphometrics is typically applied to dead specimens (museum specimens, field collections preserved in fixative) or if done using live animals, the method of image acquisition for two-dimensional (2D) analysis is often photography.

Dalton et al. (2017) investigated the effects of age, sex, and beak size on beak shape variation in domestic turkeys using photography and geometric morphometrics. The authors reported a wide range of phenotypic variation in turkey beak shape with sex and beak size having a much greater effect on shape than age. The shape patterns reported by Dalton et al. (2017) are similar to those found in other bird species (Foster et al., 2008; Kulemeyer et al., 2009; Shao et al., 2016). Although they did not look at the beak specifically, Stange et al. (2018) compared skull morphology between domesticated chicken breeds and wild fowl using geometric morphometrics analysis of microscribe-acquired landmark coordinates. Domesticated chickens occupied a greater portion of morphospace (shape space) and therefore had greater variation in skull shape compared to their wild fowl ancestors (Stange et al., 2018).

The objective of this study was to radiograph breeding stock consisting of two different pure lines of White Leghorn hens to 1) generate bone morphometric data and 2) describe the phenotypic variation in premaxillary (within the top beak) and dentary (within the bottom beak) bone shape that exists within these two populations. The collection of these phenotypes will allow for the calculation of genetic parameters (heritability, QTL, and estimated breeding values) in future studies. As mentioned, geometric morphometrics is typically applied to dead specimens or if using live specimens, applied to photographs. To our knowledge, geometric morphometrics has not been applied to radiographs for the assessment and characterization of beak morphology in poultry. Radiography offers the advantage of being able to analyze the bones, which may contribute to overall beak shape and

size. This study helped develop the novel capability of using radiography for the application of geometric morphometrics in live, non-sedated laying hens.

MATERIALS AND METHODS

This study was reviewed and approved by the Animal Welfare and Ethical Review Body at the Roslin Institute (University of Edinburgh). This study was conducted in the UK under a Home Office project license (70/7909) and complied with UK regulations regarding the treatment of experimental animals (Home Office (UK), 2014).

Animals and Housing

Birds ($n = 825$) used for this study were from two pure line populations (Line A and B) of White Leghorn laying hens. Both Line A ($n = 387$) and Line B ($n=432$) hens contribute to the four-line breeding scheme of Lohmann Selected Leghorn (LSL) hybrid layers (Lohmann Breeders GmbH). Hens from both populations were housed in conventional cages ($n = 200$) in family groups (full-siblings and/or half-siblings) with a maximum of five birds per cage.

Radiographic Examination

Equipment. The radiograph device used was a MeX+20BT Lite portable x-ray generator with a Cuattro Slate 6DR system (Figure 1a). The generator was suspended over the radiograph plate and a 2-metre controlled radiation zone was demarcated using plastic chain-link barriers. A lead curtain separated the controlled radiation zone from the bird handling zone.

To reduce handlers' exposure to radiation, custom restraint devices were developed to hold the hens in place. The restraint was performed using methods adapted for veterinary

radiograph examination. A restraint jacket was made of cotton denim. Birds were wrapped in the cotton jacket, which was then secured by an attached strap of hook and loop tape (Figure 1b). A separate strap of hook and loop tape was used to secure the legs together. For immobilizing the head, a cotton hood was custom designed (Figure 1c). The hood could be placed over the head and the beak would show through a small opening at one corner. The hood was secured underneath the head by small hook and loop fasteners. These could be adjusted to accommodate different sized heads and combs. Once the hen was placed in right lateral recumbence on the radiograph plate, the cloth hood was secured to a strap of hook and loop tape running along the radiograph plate. For optimal positioning, a small piece of radiolucent foam was placed underneath the beak for the duration of the radiograph.

Procedure. All hens from two cages were removed from their home cage, loaded into an enclosed transport trolley, and transported to the radiation zone (up to 10 hens per transport trolley). Each side of the trolley had five smaller cages in which one hen was placed and each home cage took up one side of the trolley. Hens were removed from the trolley one at a time, had their wing band recorded, and then were restrained as described above. The handlers left the radiation zone and a lateral radiograph was taken. The hen remained conscious and non-sedated for the entire procedure. The radiograph was checked for image quality (e.g., blurriness). If the radiograph was not usable, another was taken. After verification of a usable radiograph, the hen was lifted off the radiograph plate, the fabric hood and jacket were removed, and the hen was immediately returned to the trolley where it remained until the procedure was complete for all hens in the trolley. When returning the hens to their home cage, the wing band numbers belonging to each cage were verified by matching them to a sticker on the front of the cage. Following catching and placement in the transport trolley, the entire procedure lasted approximately 120 s for each hen, beginning

with removal from the trolley, restraint, radiograph acquisition, and return to the trolley. A total of 825 radiographs (one per hen) were collected over nine days.

Radiograph Analysis

Formatting. Left lateral radiographs (DICOM format) were cropped to 600 x 600 pixels and uniformly sharpened (Sharpen feature) using ImageJ analysis software (v.1.53g; National Institutes of Health, Bethesda, MD, USA). Radiographs were cropped to show the entire head and the C1 vertebra. Radiographs were excluded from landmarking if the image was blurry or if the landmarks could not be properly placed on their locations. This resulted in 819 and 710 lateral radiographs being available for premaxillary and dentary landmarking, respectively. The difference in the number of radiographs available for premaxillary vs. dentary landmarking was primarily due to blurriness from the hen moving her lower mandible during radiograph acquisition.

Placement of landmark coordinates. The radiographs were landmarked in ImageJ using the Multi-point Tool. The anatomical terms of location used to describe the placement of the landmarks and the shape of both bones are provided in Figure 2. Thirteen landmarks (LMs) were chosen for the premaxillary and dentary bones (Figure 3). LMs 1 to 3 were placed on locations that had the same relative position (homology) between specimens (Zelditch et al., 2012). LMs 4 to 13 consisted of sliding semilandmarks and helped quantify the shape of the bone (Gunz and Mitteroecker, 2013). These LMs slid along the curves of the bones until their positions were optimized to minimize shape differences (Gunz and Mitteroecker, 2013). The 2D landmark coordinates were copied from the Results table in ImageJ into a Microsoft Excel spreadsheet. The spreadsheet (.xlsx format) was imported into R (v.3.6.1) and then R-based scripting and data visualization were done using the integrated development environment RStudio (v.1.2.5001) (R Core Team, 2019).

Geometric morphometric analysis. Multivariate shape analysis was done using the R package Geomorph (v.4.0.0) (Adams et al., 2021). Outliers in the upper quartile as shown by Procrustes distance to the mean (square root of the sum of squared distances between the individual shape measurement and the consensus shape) were removed from analysis; this reduced the dataset of 2D landmark coordinates to 806 premaxillary and 684 dentary lateral radiographs. Using Procrustes superimposition, the 2D landmark coordinates for all the radiographs (each bone separately) were standardized such that landmarks were superimposed, re-scaled, and rotated to a common orientation. The resulting transformed landmarks are termed Procrustes coordinates. A distance matrix of individuals was calculated from the Procrustes coordinates. The covariation described by the matrix was decomposed by principal components analysis (PCA) and eigenvectors and their corresponding eigenvalues were found. The eigenvector with the highest eigenvalue was the first principal component (PC) and explained the most covariation in shape. Each sequential PC (eigenvector) described continuously smaller covariation (eigenvalue). These eigenvalues allowed the shape variables to be ordered in morphospace. From there, similarities and differences in shape were interpreted from where each individual point (corresponding to one hen) was on the PCA plot.

The Procrustes superimposition also created a consensus bone shape by identifying the centroid (center point) of all the landmarks and semilandmarks in each radiograph. The centroid size was then calculated as the square root of the sum of squared distances of landmarks from the centroid (Zelditch et al., 2012). In the present study, centroid size served as a proxy for relative bone size.

Statistical analysis. To test for allometry (the influence of size on shape) for each bone, a multivariate regression of the Procrustes shape coordinates onto log centroid size for

the entire dataset was performed. A Procrustes ANOVA was performed to determine if bone shape variation and allometry differed between the two pure lines. Data were analyzed using the `procD.lm` function within Geomorph. Significance was evaluated with a residual randomization permutation procedure with 1000 iterations. A Spearman correlation using the `cor.test` function in the R Stats package was performed to test the relationship between 1) PC1 scores of the premaxillary and dentary bones (i.e., the relationship between the two bone shapes) and 2) log centroid sizes of the premaxillary and dentary bones (i.e., the relationship between the two bone sizes). Differences were considered significant when $P \leq 0.05$.

RESULTS

Premaxillary Bone

For the premaxillary bone, there were three principal components that explained the majority of the total variation in premaxillary bone shape in the two populations of pure line hens. PC1 accounted for 56% of the total variation. Though the two lines overlapped in morphospace, there was noticeable separation between them. In contrast to Line A, Line B tended towards premaxillary shapes with more pronounced downward curvature. Also, the caudal end of the premaxillary bone tended to be deeper in Line B (Figure 4). PC2 accounted for 18% of the total shape variation and describes a rostral (PC2 min) vs. caudal (PC2 max) shift in the dorsal and ventral margins of the bone and a corresponding caudal (PC2 min) vs. rostral (PC2 max) shift of the bone tip (Figure 4). PC3 explained 11% of the total shape variation and described a rostral (PC3 min) vs. caudal (PC3 max) shift in the dorsal/ventral beak margins as well as a slight narrowing (PC3 min) vs. widening (PC3 max) of the caudal-most ventral margins of the premaxillary bone (Figure 5).

To test whether there was a relationship between bone size and its shape (static allometry), the Procrustes shape coordinates for the entire dataset were regressed on log centroid size. The regression revealed that the shape coordinates were significantly associated with bone size (Z-score = 10, $P < 0.01$) and that bone size accounted for 42% of the total shape variation. Based on Procrustes ANOVA, premaxillary bone shape also differed significantly between the two lines (Z-score = 8, $P < 0.01$). Allometry was also unique between the lines as the interaction between log centroid size and line was significant ($P = 0.04$). Line A hens had larger log centroid sizes (mean = 6.11 ± 0.002) compared to Line B (mean = 6.04 ± 0.003). Because this interaction was significant, no attempt was made to correct for allometry.

Dentary Bone

The PCA of the dentary bone revealed two principal components that explained 81% of the total shape variation in the two lines. PC1 accounted for 62% of the shape variation and described a widening and lengthening (PC1 min) vs. a narrowing and shortening (PC1 max) of the bone (Figure 6). As observed with the premaxillary bone, PC1 appeared to separate the lines, with a longer/wider dentary bone more characteristic of Line B. PC2 accounted for 19% of the total variation and described the angle between the dentary bone and its articular process associated with an inferior (PC2 min) vs. superior (PC2 max) shift of the caudal end of the dentary bone and a superior (PC2 min) vs. inferior (PC2 max) shift of the bone tip (Figure 6).

Multivariate regression of the Procrustes shape coordinates on log centroid size for the entire dataset showed that dentary bone shape was significantly associated with its size (Z-score = 8, $P < 0.01$) and that bone size accounted for 42% of the total dentary shape variation. Like the premaxillary bone, dentary bone shape differed significantly between lines

(Z-score = 7, $P < 0.01$) as did allometry ($P < 0.01$). Line A had larger log centroid sizes (mean = 5.81 ± 0.005) compared to Line B (mean = 5.68 ± 0.007). As with the premaxillary bone, the interaction between log centroid size and line was significant for the dentary bone ($P < 0.01$) so no attempt to correct for allometry was made.

Spearman Correlation

Correlation between the PC1 scores of the premaxillary and dentary bones found that there was a strong, positive relationship ($r_s = 0.69$; $P < 0.01$) between the shapes of the premaxillary and dentary bones. Using log centroid size as a proxy for bone size, a strong, positive relationship between the sizes of the premaxillary and dentary bones was also found ($r_s = 0.96$, $P < 0.01$).

DISCUSSION

Many previous studies examining skull and beak morphology in different avian species have done so from an evolutionary and developmental perspective (Foster et al., 2008; Bright et al., 2016, 2019; Shao et al., 2016; Cheng et al., 2017), focusing on how the beak and skull have morphologically evolved in response to each species' particular ecological niche. This differs from the present study whose aim is to understand the underlying shape of the bones within the beak so that it potentially can be used as a tool to help improve management of non-beak treated laying hen flocks. To our knowledge, this is the first study to analyze and describe the morphology of the bones within chicken beaks using radiography and landmark-based geometric morphometrics. As mentioned previously, geometric morphometric techniques are typically used on dead specimens. A challenge to collecting beak morphology data (phenotypes) for use in further genomic and behavioral studies is collecting it in live animals. Geometric morphometrics has been applied to

photographs in live turkeys (Dalton et al., 2017); however, photography does not allow for the analysis of bone shape. Radiography offers the advantage of being able to analyze both beak and bone shape but there are technical challenges with collecting radiograph data from live, non-sedated birds as birds need to be restrained during x-ray capture to limit bird and human exposure to radiation. This study showed that meaningful phenotype data can be collected from live laying hens using the methods described.

The present study found that premaxillary bone shape variability explained by PC1 and PC2 ranged from long, narrow bones with pointed tips to short, wide bones with more curved tips. Variability in dentary bone shape explained by PC1 and PC2 ranged from short and narrow to long and wide with corresponding superior vs. inferior shifts of the bone tip. The closest related study to the present one would be that of Dalton et al. (2017) who examined the external beak shape of domestic turkeys. The authors found that top beak shape in domestic turkeys ranged from short, narrow beaks with short, pointed beak tips to long, wide beaks with long, curved beak tips. Bottom beak shape in turkeys ranged from wide and round to narrow and thin with corresponding shifts in the beak tip (Dalton et al., 2017).

The axes described by both PC1 and PC2 for premaxillary bone shape variation in this study are similar to top beak shapes reported previously in other avian species: long and narrow vs. short and wide (Foster et al., 2008; Kulemeyer et al., 2009; Shao et al., 2016; Dalton et al., 2017). Although it is presumed that beak shape reflects the underlying structures of the premaxillary and dentary bones, we are unaware of any study that has formally tested this. Thus, caution is required when comparing our results to those whose analyses are based on beak shape. Regarding the curvature of the premaxillary bone tip, the present study found that hens with long and narrow premaxillary bones had more pointed (less curved) bone tips while hens with short and wide premaxillary bones had more curved

(less pointed) bone tips. In the present study it is difficult to determine which bone tip shape (pointed vs. more curved) results in a sharper external beak, particularly since there is often keratin growth extending beyond the tip of the premaxillary bone which could affect the curvature. The two lines were partially separated by PC1 for the premaxillary bone, which suggests that there are distinct bone phenotypes (particularly regarding the curvature and allometry) within each genetic line. This information is important as the incorporation of multivariate shape data into selection indices could help guide the selection of hens whose beak shapes are less apt to cause damage during SFP.

There were also mild similarities in the dentary bone shapes described in the present study and the bottom beak shapes reported by Dalton et al. (2017). In both studies, shape variability described by PC1 and PC2 could be interpreted as a widening vs. narrowing of the bone or beak with corresponding superior vs. inferior shifts of the bone or beak tip. However, once again these comparisons should be made with caution. Here, bone size accounted for approximately 42% of both the premaxillary and dentary bone shape. This is similar to what has been reported in raptors (Bright et al., 2016) and for the bottom beak of domestic turkeys (Dalton et al., 2017) and suggests that bone shape is strongly influenced by bone size. However, more research investigating the relationship between bone shape and size and the influence of body size is needed. Further studies could also incorporate lateral landmarks on different locations of the skull to determine the relationship between skull and beak shape and size. Beak and skull shape in avian species are highly integrated, meaning that change in one correlates to change in the other (Bright et al., 2016; Stange et al., 2018). It is still not fully understood to what magnitude skull shape impacts beak shape and how size factors in, although Stange et al. (2018) reported a weak effect of skull size on shape in domestic chicken breeds.

The positive correlations for both the PC1 scores and the log centroid sizes across both lines suggest a coordination of the growth and subsequent shape between the bones (i.e., the dentary bone grows and is shaped in relation to the premaxillary). This coordination may be explained by the developmental origins of beak morphology. Cranial neural crest cells contain the patterning information responsible for beak morphology variation between different avian species and both the premaxillary and dentary bone are derived from these cells (Schneider and Helms, 2003). Cranial neural crest cells may regulate the development and growth of the dentary bone such that it always fits inside the premaxillary and in fact, the top beak does contain a groove that the bottom beak fits in to (also known as the bearing horn) (Lucas and Stettenheim, 1972). However, to our knowledge, this has yet to be formally verified and because of this, caution should be used when interpreting the significant correlations as biologically important.

The results of this study demonstrate significant phenotypic variation in the shape of both the premaxillary and dentary bone within two populations of pure line White Leghorn laying hens. The present study is the first in a series that will investigate the impact of beak shape on feather pecking-related damage in laying hens. The aim of this study was to identify various bone phenotypes as an initial step rather than to associate a certain phenotype with damage. Research into why hens feather peck and other methods of mitigating the behavior is ongoing and the present study helps contribute to that knowledge by laying the foundation to investigate if the beak itself can be used as a mitigation tool. Subsequent morphometric analyses of these two hen populations will analyze the external shape of the beak and help elucidate the relationship between bone and beak shape and its impact on SFP behavior, plumage cover, and mortality. These bone and beak phenotypes will also be used in quantitative genetic analyses for the calculation of beak and bone shape heritability and to identify QTL underlying beak morphology.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Lohmann Breeders and their farm staff for their technical assistance and providing access to their facilities and birds. Thank you to the National Avian Research Facility for allowing the use of their facilities for the testing of the restraint devices during the early phases of this study.

DECLARATION OF INTEREST

The authors declare that they have no competing interests except for SS who received financial support from Lohmann Breeders and BA and MS who are employees of Lohmann Breeders.

ROLE OF THE FUNDING SOURCES

SS is supported by a Research Excellence Grant at Scotland's Rural College (SRUC), the Flexible Talent Mobility Award scheme, and Lohmann Breeders. VS receives funding from the Scottish Government Rural Affairs, Food, and the Environment (RAFE) strategic research portfolio 2016-21. JJS and ICD receive BBSRC institution strategic program grant support [BBS/E/D/30002275, BBS/E/D/30002276]. Apart from Lohmann Breeders, none of the funding sources had any involvement in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to submit the article for publication. Lohmann Breeders provided access to their facilities and birds and assisted with the collection of the head radiographs. MS and BA provided comments on the manuscript.

Declaration of interests

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Adams, D. C., M. L. Collyer, A. Kaliontzopoulou, and E. K. Balken. 2021. Geomorph: Software for geometric morphometric analyses. <https://cran.r-project.org/web/packages/geomorph/index.html>.
- Bennewitz, J., S. Bögelein, P. Stratz, M. Rodehutschord, H. P. Piepho, J. B. Kjaer, and W. Bessei. 2014. Genetic parameters for feather pecking and aggressive behavior in a large F2 -cross of laying hens using generalized linear mixed models. *Poult. Sci.* 93:810–817.
- Bilčík, B., and L. J. Keeling. 2000. Relationship between feather pecking and ground pecking in laying hens and the effect of group size. *Appl. Anim. Behav. Sci.* 68:55–66.
- Bright, J. A., J. Marugán-Lobón, S. N. Cobb, and E. J. Rayfield. 2016. The shapes of bird beaks are highly controlled by nondietary factors. *Proc. Natl. Acad. Sci. U. S. A.* 113:5352–5357.
- Bright, J. A., J. Marugán-Lobón, E. J. Rayfield, and S. N. Cobb. 2019. The multifactorial nature of beak and skull shape evolution in parrots and cockatoos (Psittaciformes). *BMC Evol. Biol.* 19:104.
- Brinker, T., P. Bijma, J. Visscher, T. B. Rodenburg, and E. D. Ellen. 2014. Plumage condition in laying hens: Genetic parameters for direct and indirect effects in two purebred layer lines. *Genet. Sel. Evol.* 46:33.
- British Egg Industry Council. 2019. Industry data. Accessed May 2020. <https://www.egginfo.co.uk/egg-facts-and-figures/industry-information/data>.
- Carruthers, C., T. Gabrush, K. Schwean-Lardner, T. D. Knezacek, H. L. Classen, and C.

- Bennett. 2012. On-farm survey of beak characteristics in White Leghorns as a result of hot blade trimming or infrared beak treatment. *J. Appl. Poult. Res.* 21:645–650.
- Centner, T. J. 2010. Limitations on the confinement of food animals in the United States. *J. Agric. Environ. Ethics* 23:469–486.
- Cheng, Y., B. Gao, H. Wang, N. Han, S. Shao, S. Wu, G. Song, Y. E. Zhang, X. Zhu, X. Lu, Y. Qu, and F. Lei. 2017. Evolution of beak morphology in the Ground Tit revealed by comparative transcriptomics. *Front. Zool.* 14:1–13.
- Craig, J. V., and W. M. Muir. 1993. Selection for reduction of beak-inflicted injuries among caged hens. *Poult. Sci.* 72:411–420.
- Dalton, H. A., B. J. Wood, T. M. Widowski, M. T. Guerin, and S. Torrey. 2017. An analysis of beak shape variation in two ages of domestic turkeys (*Meleagris gallopavo*) using landmark-based geometric morphometrics. *PLoS One* 12:e0185159.
- Dennis, R. L., A. G. Fahey, and H. W. Cheng. 2009. Infrared beak treatment method compared with conventional hot-blade trimming in laying hens. *Poult. Sci.* 88:38–43.
- Department for Environment Food and Rural Affairs. 2010. The Mutilations (Permitted Procedures) (England) (Amendment) Regulations 2010.
<https://www.legislation.gov.uk/ukxi/2010/3034/contents>.
- Dunne, C., and C. Sietou. 2020. UK consumers' willingness to pay for laying hen welfare. *Br. Food J.* 122:2867–2880.
- Ellen, E. D., and P. Bijma. 2019. Can breeders solve mortality due to feather pecking in laying hens? *Poult. Sci.* 98:3431–3442.
- Ellen, E. D., M. van der Sluis, J. Siegfors, O. Guzhva, M. J. Toscano, J. Bennewitz, L. E. van der Zande, J. A. J. van der Eijk, E. N. de Haas, T. Norton, D. Piette, J. Tetens, B. de Klerk, B. Visser, and T. B. Rodenburg. 2019. Review of sensor technologies in animal

breeding: Phenotyping behaviors of laying hens to select against feather pecking.

Animals 9:108.

Foster, D. J., J. Podos, and A. P. Hendry. 2008. A geometric morphometric appraisal of beak shape in Darwin's finches. *J. Evol. Biol.* 21:263–275.

Gentle, M. J., B. O. Hughes, A. Fox, and D. Waddington. 1997. Behavioural and anatomical consequences of two beak trimming methods in 1- and 10-d-old domestic chicks. *Br. Poult. Sci.* 38:453–463.

Gentle, M. J., and D. E. F. McKeegan. 2007. Evaluation of the effects of infrared beak trimming in broiler breeder chicks. *Vet. Rec.* 160:145–148.

Grams, V., S. Bögelein, M. A. Grashorn, W. Bessei, and J. Bennewitz. 2015. Quantitative genetic analysis of traits related to fear and feather pecking in laying hens. *Behav. Genet.* 45:228–235.

Grant, B. R., and P. R. Grant. 1993. Evolution of Darwin's finches caused by a rare climatic event. *Proc. R. Soc. B Biol. Sci.* 251:111–117.

Gunz, P., and P. Mitteroecker. 2013. Semilandmarks: A method for quantifying curves and surfaces. *Hystrix* 24:103–109.

Home Office (UK). 2014. Animals (Scientific Procedures) Act 1986 Amendment Regulations 2012. Accessed Jun. 2020. <https://www.gov.uk/government/publications/consolidated-version-of-aspa-1986>.

Icken, W., D. Cavero, and M. Schmutz. 2017. Selection on beak shape to reduce feather pecking in laying hens. *Lohmann Inf.* 51:22–27.

Jendral, M. J., and F. E. Robinson. 2004. Beak trimming in chickens: Historical, economical, physiological and welfare implications, and alternatives for preventing feather pecking and cannibalistic activity. *Avian Poult. Biol. Rev.* 15:9–23.

- Kulemeyer, C., K. Asbahr, P. Gunz, S. Frahnert, and F. Bairlein. 2009. Functional morphology and integration of corvid skulls - a 3D geometric morphometric approach. *Front. Zool.* 6:1–14.
- Kuo, F. L., J. V. Craig, and W. M. Muir. 1991. Selection and beak-trimming effects on behavior, cannibalism, and short-term production traits in White Leghorn pullets. *Poult. Sci.* 70:1057–1068.
- Lay, D. C., R. M. Fulton, P. Y. Hester, D. M. Karcher, J. B. Kjaer, J. A. Mench, B. A. Mullens, R. C. Newberry, C. J. Nicol, N. P. O’Sullivan, and R. E. Porter. 2011. Hen welfare in different housing systems. *Poult. Sci.* 90:278–294.
- Leeson, S., and T. Walsh. 2004. Feathering in commercial poultry II. Factors influencing feather growth and feather loss. *Worlds. Poult. Sci. J.* 60:52–63.
- Lucas, A. M., and P. R. Stettenheim. 1972. *Avian Anatomy. Integument Part II.* Washington, D.C.
- Lunam, C. A. 2005. The anatomy and innervation of the chicken beak: effects of trimming and re-trimming. Pages 51–73 in *Poultry Welfare Issues: Beak Trimming*. Glatz, P.C., ed. Nottingham University Press, UK.
- Lunam, C. A., P. C. Glatz, and Y. Hsu. 1996. The absence of neuromas in beaks of adult hens after conservative trimming at hatch. *Aust. Vet. J.* 7:4–7.
- Marchant-Forde, R. M., A. G. Fahey, and H. W. Cheng. 2008. Environment, well-being, and behavior: Comparative effects of infrared and one-third hot-blade trimming on beak topography, behavior, and growth. *Poult. Sci.* 87:1474–1483.
- McKeegan, D. E. F., and A. W. Philbey. 2012. Chronic neurophysiological and anatomical changes associated with infrared beak treatment and their implications for laying hen welfare. *Anim. Welf.* 21:207–217.

- Muir, W. M., H. W. Cheng, and C. Croney. 2014. Methods to address poultry robustness and welfare issues through breeding and associated ethical considerations. *Front. Genet.* 5:407.
- Napolitano, F., A. Girolami, and A. Braghieri. 2010. Consumer liking and willingness to pay for high welfare animal-based products. *Trends Food Sci. Technol.* 21:537–543.
- R Core Team. 2019. R: A language and environment for statistical computing. <https://www.r-project.org/>.
- Rodenburg, T. B., P. Bijma, E. D. Ellen, R. Bergsma, S. de Vries, J. E. Bolhuis, B. Kemp, and J. A. M. van Arendonk. 2010. Breeding amiable animals? Improving farm animal welfare by including social effects in breeding programmes. *Anim. Welf.* 19:77–82.
- Rodenburg, T. B., A. J. Buitenhuis, B. Ask, K. A. Uitdehaag, P. Koene, J. J. van der Poel, and H. Bovenhuis. 2003. Heritability of feather pecking and open-field response of laying hens at two different ages. *Poult. Sci.* 82:861–867.
- Savory, C. J. 1995. Feather pecking and cannibalism. *Worlds. Poult. Sci. J.* 51:215–219.
- Schneider, R. A., and J. A. Helms. 2003. The cellular and molecular origins of beak morphology. *Science* 299:565–568.
- Scottish Executive. 2010. The Prohibited Procedures on Protected Animals (Exemptions) (Scotland) Regulations 2010. <https://www.legislation.gov.uk/ssi/2010/387/contents/made>.
- Shao, S., Q. Quan, T. Cai, G. Song, Y. Qu, and F. Lei. 2016. Evolution of body morphology and beak shape revealed by a morphometric analysis of 14 Paridae species. *Front. Zool.* 13:30.
- Spooner, J. M., C. A. Schuppli, and D. Fraser. 2014. Attitudes of Canadian citizens toward farm animal welfare: A qualitative study. *Livest. Sci.* 163:150–158.

Stange, M., D. Núñez-León, M. R. Sánchez-Villagra, P. Jensen, and L. A. B. Wilson. 2018.

Morphological variation under domestication: how variable are chickens? *R. Soc. Open Sci.* 5:180993.

Struthers, S., H. L. Classen, S. Gomis, T. G. Crowe, and K. Schween-Lardner. 2019. The

impact of beak tissue sloughing and beak shape variation on the behavior and welfare of infrared beak-treated layer pullets and hens. *Poult. Sci.* 98:4269–4281.

United Egg Producers. 2021. Facts and stats. Accessed Apr. 2021.

<https://unitedegg.com/facts-stats/>.

Zelditch, M., D. Swiderski, and H. Sheets. 2012. Geometric morphometrics for biologists.

2nd ed. Academic Press.




Figure 1. (a) Set up of the portable radiograph device on the laying hen farm. Arrow indicates radiograph plate. Body (b) and head (c) restraint devices used during radiograph acquisition.




Figure 2. Schematic demonstrating the anatomical terms of location used to describe premaxillary (within the top beak) and dentary (within the bottom beak) bone shape. The anatomical terms of location are in relation to the white shaded area on each bone.




Figure 3. The landmarks (LMs) and semilandmarks used for the analyses of the left lateral radiographs. (a) premaxillary bone (LM 1, tip of premaxillary bone; LM 2, rostral end of the nares; LM 3, base of skull where spinal cord enters; LMs 4-13, semilandmarks) and (b) dentary bone (LM 1, rostral tip of dentary bone; LM 2, caudal-most end of dentary bone; LM 3, caudal-most end of angular bone; LMs 4-13, semilandmarks).




Figure 4. The premaxillary bone shape variation explained by PC1 and PC2 for Line A (black) and B (grey) pure line hens. The point figures at each end of the x and y axes are visual representations of the premaxillary bone shape. The blue lines represent the mean shape of the premaxillary bone for the two lines. The red lines represent the premaxillary bone shape at the minimum and maximum values along the axes of PC1 and PC2.




Figure 5. The premaxillary bone shape variation explained by PC1 and PC3 for Line A (black) and B (grey) pure line hens. The point figures at each end of the x and y axes are visual representations of the premaxillary bone shape. The blue lines represent the mean shape of the premaxillary bone for the two lines. The red lines represent the premaxillary bone shape at the minimum and maximum values along the axes of PC1 and PC3.




Figure 6. The dentary bone shape variation explained by PC1 and PC2 for Line A (black) and B (grey) pure line hens. The point figures at each end of the x and y axes are visual representations of the dentary bone shape. The blue lines represent the mean shape of the dentary bone for the two lines. The red lines represent the dentary bone shape at the minimum and maximum values along the axes of PC1 and PC2.